

IRF Considerations – Tiers

C Tier

- Dysphagia
 - RIC Exclusion 01

D Tier

- NT Hemorrhage (Subarachnoid, intracerebral, Subdural)
 - RIC Exclusion: 01, 02, 03
- Cerebral Infarction
 - Ric Exclusion: 01
- **Note- Hemiplegia not associated to CVA and Flaccid are D tiers but not for stroke patients.**

Activity

74-year-old right-handed male admitted to rehab status cerebral infarction due to thrombosis of left middle cerebral artery. The provider documents patient has right sided hemiplegia, oral dysphagia, aphasia, and right sided and facial droop.

- **Code Etiologic and Comorbid Conditions**
- **Are there tiered comorbidities**
- **UB-04 codes/PDx and comorbidities**

Other AMI Options

- Diagnosed on EKG but without symptoms: (I25.2)
- Healed/Old/Past: (I25.2)
- Intraoperative
 - During cardiac surgery: (I97.790)
 - During other surgery: (I97.791)
- Post Myocardial Infarction Syndrome: (I24.1)

Activity

A patient is seen for an acute
subendocardial myocardial infarction

Activity

A patient underwent a percutaneous coronary intervention and during the procedure the patient has a type 4A Myocardial Infarction

Activity

A patient is seen diagnosed with a STEMI of the inferior wall. This STEMI is subsequent to a previous STEMI suffered 15 days prior